

**Amendments to the Claims**

Claim 1 (Currently Amended): A method for amplifying a cDNA comprising:  
obtaining an mRNA;  
reverse transcribing the mRNA into cDNA with reverse transcriptase without RNase H activity  
so that a cDNA-mRNA complex is formed;  
degrading the mRNA from the cDNA-mRNA complex to form a linear cDNA;  
ligating the ends of said linear cDNA to form a circular cDNA;  
introducing first and second sequence specific primers to said circular cDNA wherein said first primer comprises a 3' end of the same which is toward the 5' end of said circular cDNA and said second primer comprises a 3' end of the same which is toward the 3' end of said circular cDNA ; and  
initiating a primer extension amplification reaction to increase copy number of said circular cDNA.

Claim 2 (Cancelled)

Claim 3 (Original): The method of claim 1 wherein said primer extension amplification reaction is a polymerase chain reaction.

Claim 4 (Original): The method of claim 1 wherein said polymerase chain reaction is employed with Taq polymerase or other heat-resisted DNA polymerase.

Claim 5 (Original): The method of claim 1 wherein said PCR is touchdown PCR.

Claim 6 (Previously Presented): The method of claim 1 further comprising the step of:  
harvesting said amplified cDNA.

Claim 7 (Original): The method of claim 1 wherein said ligase is T4 DNA ligase.

Claim 8 (Original): The method of claim 1 wherein said primer is a degenerate primer.

Claim 9 (Previously Presented): The method of claim 1 wherein said first and second primers are designed to hybridize to from about 4 to about 35 contiguous bases from a sequence known or suspected to be present in said circular cDNA.

Claims 10-11 (Cancelled)

Claim 12 (Currently Amended): A method for amplifying a cDNA, including the 5' and 3' ends, comprising:  
obtaining an mRNA;  
contacting the mRNA with reverse transcriptase without RNase H so that a cDNA-mRNA complex is formed;  
degrading the mRNA from the cDNA-mRNA complex to form a linear cDNA;  
circularizing said cDNA  
contacting the circularized cDNA with first and second sequence specific primers wherein said first primer comprises a 3' end of the same which is toward the 5' end of said circular cDNA and said second primer comprises a 3' end of the same which is toward the 3' end of said circular cDNA; and  
introducing a polymerase and a supply of nucleotide bases to said circularized cDNA so that an amplification reaction occurs, wherein said region of said cDNA outside of said first and second primers including the 3' and 5' ends of said cDNA is amplified.

Claim 13 (Previously Presented): The method of claim 12 wherein said ligase is T4 DNA ligase.

Claim 14 (Original): The method of claim 1 wherein said primer is a degenerate primer.

Claim 15 (Previously Presented): The method of claim 1 wherein said forward and reverse primers are designed to hybridize to from about 4 to about 35 contiguous bases from a sequence known or suspected to be present in said circular cDNA.

## Claims 16-25 (Cancelled)

Claim 26 (Currently Amended): A method for amplifying a cDNA comprising:  
obtaining an mRNA;  
reverse transcribing the mRNA into cDNA with reverse transcriptase without RNase H activity  
so that a cDNA-mRNA complex is formed;  
degrading the mRNA from the cDNA-mRNA complex to form a linear cDNA;  
ligating the ends of said linear cDNA to form a circular cDNA;  
introducing first and second sequence specific promoters wherein said promoters are degenerate  
primers and wherein said first primer comprises a 3' end of the same which is toward the  
5' end of said circular cDNA and said second primer comprises a 3' end of the same  
which is toward the 3' end of said circular cDNA; and  
initiating a primer extension amplification reaction to increase copy number of said circular  
cDNA.

Claim 27 (Currently Amended): A method for amplifying a cDNA comprising:  
obtaining an mRNA;  
reverse transcribing the mRNA into cDNA with reverse transcriptase without RNase H activity  
so that a cDNA-mRNA complex is formed;  
degrading the mRNA from the cDNA complex to form a linear cDNA;  
ligating the ends of said circular cDNA to form a circular cDNA;  
introducing first and second sequence specific primers to said circular cDNA, wherein said first  
and second primers are designed to hybridize to from about 4 to about 35 contiguous  
bases from a sequence known or suspected to be present in said circular cDNA and  
wherein said first primer comprises a 3' end of the same which is toward the 5' end of said  
circular cDNA and said second primer comprises a 3' end of the same which is toward the  
3' end of said circular cDNA; and  
initiating a primer extension amplification reaction to increase copy number of said circular  
cDNA.

## Claims 28-31 (Cancelled)